

Duncan M. Tooke,* M. Lutz and
Anthony L. SpekBijvoet Center for Biomolecular Research,
Department of Crystal and Structural Chemistry,
Utrecht University, Padualaan 8, 3584 CH
Utrecht, The Netherlands

Correspondence e-mail: d.m.tooke@chem.uu.nl

Key indicators

Single-crystal X-ray study
 $T = 150$ K
Mean $\sigma(\text{C}-\text{C}) = 0.002$ Å
 R factor = 0.050
 wR factor = 0.135
Data-to-parameter ratio = 19.0For details of how these key indicators were
automatically derived from the article, see
<http://journals.iucr.org/e>.

1,3-Bis(3,5-dimethylphenoxy)propane

The molecular structure of the title compound, $\text{C}_{19}\text{H}_{24}\text{O}_2$, exhibits great similarity to that of the previously published non-derivatized molecule, the greatest difference being seen in the central torsion angles and ring orientations. The structure additionally features a π - π stacking interaction and a weak $\text{C}-\text{H}\cdots\pi$ interaction.


Received 7 June 2005

Accepted 10 June 2005

Online 17 June 2005

Comment

The title compound, (I), was characterized as a precursor for oxidation to isophthalaldehyde donor ligands, as part of an ongoing series of investigations (Zondervan *et al.*, 1997).



The aliphatic linker between the aromatic rings is not in a fully stretched conformation, but has adopted an (*anti*, *-gauche*, *-gauche*, *anti*) conformation. This is directly comparable to the published (Sasanuma *et al.*, 2004) non-derivatized ligand, *viz.* 1,3-diphenoxypropane, (II), which adopts a nearly identical conformation (Fig. 2).

The main difference between the two molecules is in the torsion angles around the bonds which share the central atom of the propane linker. In (I), these are -73.50 (15) and -72.70 (16) $^\circ$, whereas in (II) they are both -57.1 (2) $^\circ$ by symmetry. This difference in torsion angle results in a significant change in the pitch of the two sets of benzene rings. This ring orientation should not be attributed to the torsion angles





Figure 1
View of the title compound with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

Quaternion fit based on the five linking O and C atoms of the title compound (black) with the previously published non-derivatized molecule (red).

around the ring C–O bonds, which are actually quite comparable; 173.70 (13) and 177.51 (13)° in (I), and 177.2 (2)° for both rings in (II).

Although there are a number of rings in the structure of (I), there is only one instance of π – π stacking, between the C1–C6 ring and its symmetry equivalent at $(1 - x, -y, 2 - z)$. The two rings are parallel by symmetry with a perpendicular separation of 3.411 (1) Å and a horizontal slippage of 1.16 Å (Fig. 3).

There is, additionally, a weak C–H... π interaction between H11B and the C12–C17 ring system, with a hydrogen to ring centroid (C_g) distance of 2.88 Å and a C–H... C_g angle of 157° (Fig. 4).

Experimental

Single crystals, with a melting point of 308 K, were obtained by recrystallization from acetone.

Crystal data


$C_{19}H_{24}O_2$	$Z = 2$
$M_r = 284.38$	$D_x = 1.173 \text{ Mg m}^{-3}$
Triclinic, $P\bar{1}$	Mo $K\alpha$ radiation
$a = 8.7906$ (12) Å	Cell parameters from 25 reflections
$b = 9.6774$ (11) Å	$\theta = 10.0$ – 15.2°
$c = 10.7788$ (12) Å	$\mu = 0.07 \text{ mm}^{-1}$
$\alpha = 82.410$ (9)°	$T = 150$ (2) K
$\beta = 78.47$ (1)°	Block, colourless
$\gamma = 63.796$ (9)°	$0.5 \times 0.5 \times 0.4 \text{ mm}$
$V = 805.11$ (17) Å ³	

Data collection


Nonius CAD-4-Turbo diffractometer	$\theta_{\text{max}} = 27.5^\circ$
ω scans	$h = -11 \rightarrow 11$
Absorption correction: none	$k = -12 \rightarrow 12$
4628 measured reflections	$l = -13 \rightarrow 9$
3689 independent reflections	3 standard reflections
2710 reflections with $I > 2\sigma(I)$	frequency: 60 min
$R_{\text{int}} = 0.042$	intensity decay: 0.8%

Refinement

Refinement on F^2	$w = 1/[\sigma^2(F_o^2) + (0.062P)^2 + 0.2118P]$
$R[F^2 > 2\sigma(F^2)] = 0.050$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.135$	$(\Delta/\sigma)_{\text{max}} = 0.001$
$S = 1.03$	$\Delta\rho_{\text{max}} = 0.21 \text{ e \AA}^{-3}$
3689 reflections	$\Delta\rho_{\text{min}} = -0.23 \text{ e \AA}^{-3}$
194 parameters	
H-atom parameters constrained	

**Figure 3**

π – π stacking of the C1–C6 ring. The view is perpendicular to the ring mean plane.

**Figure 4**

Weak C–H... π interaction (dashed lines) between H11B and the C12–C17 ring.

Table 1

Selected geometric parameters (Å, °).

O1–C1	1.3673 (18)	O2–C11	1.4397 (18)
O1–C9	1.4351 (18)	C9–C10	1.508 (2)
O2–C12	1.3647 (18)	C10–C11	1.509 (2)
C9–O1–C1–C2	173.70 (13)	C12–O2–C11–C10	–179.40 (13)
C1–O1–C9–C10	179.91 (12)	C9–C10–C11–O2	–72.70 (16)
O1–C9–C10–C11	–73.50 (15)	C11–O2–C12–C13	177.51 (13)

All H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for methyl H atoms and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for all other H atoms.

Data collection: locally modified *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *SET4* (de Boer & Duisenberg, 1984); data reduction: *HELENA* (Spek, 1997); program(s) used to solve structure: *SHELXS96* (Sheldrick, 1996); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *PLATON* (Spek, 2003); software used to prepare material for publication: *PLATON*.

The authors thank the Council for the Chemical Sciences of the Netherlands Organization for Scientific Research (CW–NWO) for their support. C. Zondervan and B. L. Feringa (University of Groningen) are thanked for their kind donation of the sample.

References

- Boer, J. L. de & Duisenberg, A. J. M. (1984). *Acta Cryst.* A40, C–410.
Enraf–Nonius (1989). *CAD-4 Software*. Version 5. Enraf–Nonius, Delft, The Netherlands.

- Sasanuma, Y., Ono, T., Kuroda, Y., Miyazaki, E., Hikino, K., Arou, J., Nakata, K., Inaba, H., Tozaki, K., Hayashi, H. & Yamaguchi, K. (2004). *J. Phys. Chem. B*, **108**, 13163–13176.
- Sheldrick, G. M. (1996). *SHELXS96*. University of Göttingen, Germany.
- Sheldrick, G. M. (1997). *SHELXL97*. University of Göttingen, Germany.
- Spek, A. L. (1997). *HELENA*. Utrecht University, The Netherlands.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Zondervan, C., van den Beuken, E. K., Kooijman, H., Spek, A. L. & Feringa, B. L. (1997). *Tetrahedron Lett.* **38**, 3111–3114.